

Area Between a Curve and the y-Axis

Q1, (OCR 4722, Jun 2008, Q5)

(i) $\int xdy = \int ((y-3)^2 - 2)dy$
 $= \int (y^2 - 6y + 7)dy$ **A.G.**
 $3 + \sqrt{(2+2)} = 5, \quad 3 + \sqrt{(14+2)} = 7$

- B1** Show $x = y^2 - 6y + 7$ convincingly
- B1** State or imply that required area = $\int xdy$
- B1** Use $x = 2, 14$ to show new limits of $y = 5, 7$

3

(ii) $\left[\frac{1}{3}y^3 - 3y^2 + 7y \right]_5^7$
 term
 $= \left(\frac{343}{3} - 147 + 49 \right) - \left(\frac{125}{3} - 75 + 35 \right)$
 $= 16\frac{1}{3} - 1\frac{2}{3}$
 $= 14\frac{2}{3}$

- M1** Integration attempt, with at least one correct
- A1** All three terms correct
- M1** Attempt $F(7) - F(5)$
- A1** Obtain $14\frac{2}{3}$, or exact equiv

4

Q2, (OCR 4722, Jun 2011, Q4)

<p>(i) $x + 4 = (y + 1)^2$ $x + 4 = y^2 + 2y + 1$ $x = y^2 + 2y - 3$ A.G.</p>	<p>M1 Attempt to make x the subject</p>	<p>Allow M1 for $x = (y \pm 1)^2 \pm 4$ only. Allow M1 if $(y + 1)^2$ becomes $y^2 + 1$, but only if clearly attempting to square the entire bracket – squaring term by term is M0. Must be from correct algebra, so M0 if eg $\sqrt{(x + 4)} = \sqrt{x} + \sqrt{4}$ is used.</p>
	<p>A1 2 Verify $x = y^2 + 2y - 3$</p>	<p>Need to see an extra step from $(y + 1)^2 - 4$ to given answer ie explicit expansion of bracket. No errors seen. SR B1 for verification, using $y = -1 + \sqrt{(y^2 + 2y - 3 + 4)}$, and confirming relationship convincingly, or for rearranging $x = f(y)$ to obtain given $y = f(x)$.</p>
<p>ii) $\int_1^3 (y^2 + 2y - 3) dy = \left[\frac{1}{3}y^3 + y^2 - 3y \right]_1^3$ $= (9 + 9 - 9) - (\frac{1}{3} + 1 - 3)$ $= (9) - (-1\frac{2}{3})$ $= 10\frac{2}{3}$</p>	<p>B1 State or imply that the required area is given by $\int_1^3 (y^2 + 2y - 3) dy$</p>	<p>No further work required beyond stating this. Allow if $3x$ appears in integral. Any further consideration of other areas is B0.</p>
	<p>M1 Attempt integration</p>	<p>Increase in power of y by 1 for at least two of the three terms. Can still get M1 if the -3 disappears, or becomes $3x$. Allow M1 for integrating a function of y that is no longer the given one, eg subtracted from 3, or using their incorrect rearrangement from part (i).</p>
	<p>A1ft Obtain at least two correct terms</p>	<p>Allow for unsimplified coefficients. Allow follow-through on any function of y as long as at least 2 terms and related to the area required. Condone \int, dy or $+ c$ present.</p>
	<p>M1 Attempt $F(3) - F(1)$ for their integral</p>	<p>Must be correct order and subtraction. This is independent of first M1 so can be given for substituting into any expression other than $y^2 + 2y - 3$, including $2y + 2$. If last term is $3x$ allow M1 for using 3 and 1 throughout integral, but M0 if x value is used instead.</p>
	<p>A1 5 Obtain $10\frac{2}{3}$ aef</p>	<p>Must be an exact equiv so $10.\dot{6}$ is fine (but $9\frac{5}{3}$ is A0). 10.7, $10.66\dots$ or $10\frac{2}{3} + c$ are A0. Must come from correct integral, so A0 if from $3x$. Must be given as final answer, so further work eg subtracting another area is A0 rather than ISW.</p>
	<p>7</p>	<p>Answer only is 0/5, as no evidence is provided of integration. SR Finding the shaded area by direct integration with respect to x (ie a C3 technique) can have 5 if done correctly, 4 if non-exact decimal given as final answer but no other partial credit.</p>

Q3, (OCR 4722, Jun 2014, Q9)

(i)	$0.5 \times 2.5 \times (1 + 2(-3 + 2\sqrt{6.5}) + 3)$ $= 10.2$	<p>M1*</p> <p>M1d*</p> <p>A1</p> <p>[3]</p>	<p>Attempt y-values at $x = 0, 2.5, 5$ only</p> <p>Attempt correct trapezium rule, inc $h = 2.5$</p> <p>Obtain 10.2, or better</p>	<p>M0 if additional y-values found, unless not used y_1 can be exact or decimal (2.1 or better) Allow M1 for using incorrect function as long as still clearly y-values that are intended to be the original function eg $-3 + 2\sqrt{x+4}$ (from $\sqrt{(x+4)} = \sqrt{x} + \sqrt{4}$)</p> <p>Fully correct structure reqd, including placing of y-values The 'big brackets' must be seen, or implied by later working Could be implied by stating general rule in terms of y_0 etc, as long as these have been attempted elsewhere and clearly labelled Using x-values is M0 Can give M1, even if error in evaluating y-values as long correct intention is clear</p> <p>Allow answers in the range [10.24, 10.25] if >3sf A0 if exact surd value given as final answer</p> <p>Answer only is 0/3 Using 2 separate trapezia can get full marks Using anything other than 2 strips of width 2.5 is M0 Using the trapezium rule on result of an integration attempt is 0/3</p>
(ii)	$(5 \times 3) - 10.2 = 4.8$	<p>M1</p> <p>A1FT</p> <p>[2]</p>	<p>Attempt area of rectangle – their (i)</p> <p>Obtain 4.8, or better</p>	<p>As long as $0 < \text{their (i)} < 15$</p> <p>Allow for exact surd value as well Allow answers in range [4.75, 4.80] if > 2sf</p>

(iii) $x = \frac{1}{4}(y^2 + 6y - 7)$

$$\text{area} = \left[\frac{1}{12}y^3 + \frac{3}{4}y^2 - \frac{7}{4}y \right]_1^3$$

M1	Attempt to write as $x = f(y)$	Must be correct order of operations, but allow slip with inverse operations eg $+/-$, and omitting to square the $\frac{1}{2}$ Allow $y^2 + 9$ from an attempt to square $y + 3$, even if $(y + 3)^2$ is not seen explicitly first Allow maximum of 1 error
A1	Obtain $x = \frac{1}{4}(y^2 + 6y - 7)$ aef	Allow A1 as soon as any correct equation seen in format $x = f(y)$, eg $x = \frac{1}{4}(y + 3)^2 - 4$ or $x = \frac{1}{4}(y^2 + 6y + 9) - 4$, and isw subsequent error
M1*	Attempt integration of $f(y)$	Expand bracket and increase in power by 1 for at least two terms (allow if constant term disappears) Independent of rearrangement attempt so M0M1 is possible Can gain M1 if their $f(y)$ has only two terms, as long as both increase in power by 1 Allow M1 for $k(y + 3)^3$, any numerical k , as the integral of $(y + 3)^2$ or M1 for $k(\frac{1}{2}(y + 3))^3$ from $(\frac{1}{2}(y + 3))^2$ oe if their power is other than 2
A1	Obtain $\frac{1}{12}y^3 + \frac{3}{4}y^2 - \frac{7}{4}y$ aef	Or $\frac{1}{12}(y + 3)^3 - 4y$ A0 if constant term becomes $-\frac{7}{4}x$ not $-\frac{7}{4}y$
B1	State or imply limits are $y = 1, 3$	Stated, or just used as limits in definite integral Allow B1 even if limits used incorrectly (eg wrong order, or addition) Allow B1 even if constant term is $-\frac{7}{4}x$ (or their cx)

$$= \frac{15}{4} - \left(-\frac{11}{12}\right)$$

$$= \frac{14}{3} \quad \mathbf{AG}$$

M1d* Attempt correct use of limits

A1 Obtain $\frac{14}{3}$

[7]

Correct order and subtraction
 Allow M1 (BOD) if y limits used in $-\frac{7}{4}x$ (or their cx), but
 M0 if $x=0, 5$ used
 Minimum of two terms in y
 Only term allowed in x is their c becoming cx
 Allow processing errors eg $(\frac{1}{12} \times 3)^3$ for $\frac{1}{12} \times 3^3$
 Answer is given so M0 if $\frac{14}{3}$ appears with no evidence of
 use of limits
 Minimum working required is $\frac{15}{4} + \frac{11}{12}$
 Allow M1 if using decimals (0.92 or better for $\frac{11}{12}$)
 M0 if using lower limit as $y=0$, even if $y=3$ is also used
 Limits must be from attempt at y -values, so M0 if using 0
 and 5

 Must come from exact working ie fractions or recurring
 decimals - correct notation required so A0 for 0.9166...
 A0 if $-\frac{7}{4}x$ seen in solution

SR for candidates who find the exact area by first
 integrating onto the x -axis:
B4 obtain area between curve and x -axis as $10\frac{1}{3}$
B1 subtract from 15 to obtain $\frac{14}{3}$
 And, if seen in the solution, **M1A1** for $x=f(y)$ as above